

Stafal

Antistaphylococcal Phage Lysate for Topical Application



COMPOSITON:

1 ml of the preparation contains:

Medicinal substances:

Fagi particulae antistaphylococci polyvalens, titer min. 10^7 IU in 1 ml of the solution

Auxiliary substances (adjuvants):

Thiomersal, water for injections

INDICATION GROUP:

Immunopreparation

CHARACTERISTICS:

STAFAL is an antistaphylococcal phage lysate for topical application, containing highly effective virulent phage particles with a strong and rapid lytic and polyvalent effect. The preparation is standardized in its efficacy according to the concentration of specific phage particles per 1 ml.

INDICATIONS:

STAFAL is designed exclusively for topical application in infections caused by staphylococcal strains. It can be used both in human as well as in veterinary medicine in all forms of staphylococcal infections. It is used for the destruction of staphylococcal cells in the site of progressing infection. The preparation is administered mainly for the elimination of causative agents of staphylococcal infection in the foci of infections (e.g. purulent processes of the skin, subcutis and in skin adnex) as well as in potential reservoirs (particularly in nasopharynx, intestinal and urinary tract). The preparation presents a significant therapeutical agent in the complex treatment of chronic form of staphylococcal infections (purulent affections, abscesses, fistulae, infections affecting deeply located soft tissues). STAFAL is also an important part of preventive measures in pre-operation preparation with the aim of preventing the occurrence of superponed pyogenic complications after operation interventions.

CONTRAINDICATIONS:

Hypersensitivity to some components of the preparation.

ADVERSE EFFECTS:

After the administration of preparation tingle may exceptionally occur in the site of applications.

INTERACTIONS:

No interactions have been observed.

PREGNANCY AND BREAST FEEDING:

During pregnancy and breast feeding the administration of the preparation is at the attending physician's discretion. No adverse effect is assumed in the topical application of the preparation.

DOSAGE AND MODE OF APPLICATION:

The preparation is applied into affected foci, operation wounds and potential reservoirs of infections by lavaging, compressing, spraying, by the application of tampons steeped in STAFAL. It is commonly applied twice daily according to a clinical finding. If administered into the nasal orifice, 1 - 2 drops are applied.

PRECAUTIONS:

Do not use the preparation if turbid !

STORAGE:

At a temperature + 2 °C to + 8 °C, store the inner container in the box !

WARNING:

Do not use the preparation after expiry date stated on the cover !
Keep out of reach of children !

PACKING:

10 ml in a bottle

SHELF LIFE:

1 year

Once opened, keep it at a temperature of + 2 °C to + 8 °C and use within 28 days.



Sevapharma, a.s., Průmyslová 1472/11, 102 19 Praha 10, Czech Republic
phone.: +420 296 584 631, +420 296 584 637, fax: +420 267 317 354
obchod@sevapharma.cz
www.sevapharma.cz